



# TRINITY MEDICAL MULTI-SPECIALTY GROUP

---

---

## Upper Endoscopy (EGD) and/or Endoscopic Ultrasound (EUS) prep Instructions

---

Your procedure has been scheduled for: \_\_\_\_\_

Check in time: \_\_\_\_\_

Procedure time \_\_\_\_\_

**You must cancel or reschedule 7 (seven) working days in advance in order to avoid a cancellation/reschedule fee. IF YOU DO NOT CANCEL/RESCHEDULE BEFORE PROCEDURE DATE YOU MAY BE CHARGED UP TO A \$200 CANCELLATION FEE. PHONE NUMBER FOR CANCELLING OR RESCHEDULING: (951) 501-4200.**

You will be receiving a call from our scheduler to schedule your procedure.

### **What to bring**

1. Insurance card, ID card and credit card or other form of payment if your insurance requires you to pay a co-pay.
2. Completed forms (if applicable) for the endoscopy center.
3. A list of all your prescription and over the counter medications
4. A responsible adult to take you home after the procedure
5. Warm socks for the procedure and flat shoes for after the procedure
6. Please avoid wearing contact lenses or bring jewelry/valuables

### **One week before the procedure:**

1. **STOP** using **iron**, herbals and any fish oil supplements. Preferably we also advise you to stop any non-steroidal anti-inflammatory medications like ibuprofen, motrin, aleve, naprosyn for 5 days prior to your procedure. Celebrex and Tylenol are okay.
2. If you take **Coumadin, Lovenox, Plavix, Effient, Brillinta, Ticlid, Persatine** or any medication that thins your blood please follow up with us as you may need to stop these medications 3-7 days before your procedure.
  - **Lovenox (Enoxaparin) - 12 hours prior to procedure**
  - **Aggrenox (Aspirin-Dipyridamole)- 48 hours prior to procedure**
  - **Eliquis (Apaxiban)- 48 hours prior to procedure**
  - **Pradaxa ( Dabigatran Etexilate) - 48 hours prior to procedure**
  - **Xarelto (Rivaroxaban) - 48 Hours prior to procedure**
  - **Warfarin (Coumadin)- 3 days prior to procedure**
  - **Brillianta (Ticagrelor) - 5 days prior to procedure**
  - **Effient (Prasugrel) – 7 days prior to procedure**
  - **Plavix (Clopidogrel)- 7 Days prior to procedure**
  - **Aspirin- you do not need to stop.**

**3. IF YOU TAKE GLP-1 AGONIST MEDICATIONS FOR TYPE 2 DIABETES OR WEIGHT LOSS, YOU MUST HOLD THESE MEDICATIONS FOR ONE FULL WEEK PRIOR TO YOUR PROCEDURE DATE.**

**GLP-1 AGONIST MEDICATIONS:**

- Ozempic
- Wegovy
- Mounjaro
- Trulicity
- Rybelsus
- Victoza
- Byetta
- Saxenda

**1. Diabetics please follow below:**

- **Lantus or Long Acting Insulin:** Take only ½ the dose at **midnight before** your procedure. Bring insulin in with you day of procedure
- **Glimperide (Amaryl):** Hold day of procedure
- **Glucotrol (Glipizide):** Hold day of procedure
- **Glucophage (Metformin):** Hold day of procedure
- **Actos (Pioglitazone):** Hold day of procedure
- **Invokana (Canaglifozin):** Hold day of procedure
- **Januvia (Sitagliptin):** Hold day of procedure

**Prep Instructions:**

- **MIDNIGHT:**
  - Do not have anything to eat or drink after midnight. You may take any necessary medications with a small sip of water only.
- **PROCEDURE DAY:**
  - Do not have anything to eat or drink until after your procedure is completed

By signing this form I am confirming that I have received my prep and I fully understand instructions given to me by the medical assistant.

Patient Signature: \_\_\_\_\_ Date: \_\_\_\_\_