



Trinity Medical Multi-specialty Group

SUFLAVE BOWEL PREP



WHEN YOU PURCHASE THE SUFLAVE PRESCRIPTION-- DISREGARD THE INSTRUCTIONS IN THE BOX AND FOLLOW THESE INSTRUCTIONS ONLY!

TIME: You will receive a call from the Surgery Center 2-7 days prior to your appointment with your arrival time.

You are scheduled for the day listed above, so please be sure to follow your preparation and all of the instructions detailed below. If you have access to your patient portal, the time you see listed on the schedule is NOT your scheduled time.

ARRANGE FOR A DRIVER: YOU WILL NEED A RESPONSIBLE ADULT TO DRIVE YOU HOME AFTER YOUR PROCEDURE. You will not be allowed to drive yourself home due to the sedation that you will be given. Small children are not allowed in the surgery center waiting room and drivers will be asked to wait in their car.

If you find it necessary to change or cancel your appointment, you must do this at least 7 working days prior to your appointment or you will be charged a **\$200** cancellation fee. Please call our office Monday – Friday, 8AM to 45PM at 951-517-0906 if you need to change or cancel your appointment. Follow the prompts for First Surgery Center scheduling.

Your procedure will be performed at:

- () **First Surgery Center**
31625 De Portola Rd. Suite #102
Temecula, CA 92592
Phone: (951) 517-0906

THE ITEMS THAT YOU NEED TO PURCHASE AT LEAST 72 HOURS PRIOR TO YOUR PROCEDURE:

- **SUFLAVE (This is a prescription that has been sent to your pharmacy) This is a split-dose regimen. You are required to complete both bottles.**
- **CLEAR LIQUIDS (SEE SUGGESTIONS BELOW)**

- **2 Fleet Enemas (you will bring these with you to your appointment)**

Clear Liquids you are allowed to have:

*****NO DAIRY*****

Water
 Broth (Chicken, Vegetable or Bone)
 7up, Ginger ale, Sprite
 Coffee & Tea (No cream)

*****NOTHING RED OR PURPLE*****

Apple Juice or WHITE grape juice
 Popsicles (No Red or Purple)
 Jell-O (No Red or Purple)
 Sugar or Honey

Hard Candy

ONE WEEK BEFORE YOUR PROCEDURE:

- If you are taking Blood Thinners, GLP-1, or Diabetic Medication, REVIEW AND THEN FOLLOW THE DIRECTIONS ACCORDINGLY ON THE LIST PROVIDED ON THE FOLLOWING PAGE. If you have questions regarding this, please call our office.
- Arrange for a driver. You will not be allowed to drive yourself home. Your driver must be a responsible adult. Drivers will be asked to wait in their car. We do not allow small children in the office. We will call your driver when you are ready to be picked up.
- If you find it necessary to change or cancel your appointment, you must do this at least 7 working days prior to your appointment or you will be charged a **\$200** cancellation fee. Please call our office Monday – Friday, 8AM to 4PM at 951-517-0906 if you need to change or cancel your appointment. Follow the prompts for First Surgery Center scheduling.

THE DAY BEFORE YOUR PROCEDURE: DOSE 1

YOU WILL NEED A RESPONSIBLE ADULT TO DRIVE YOU HOME AFTER YOUR PROCEDURE. Confirm with your driver, as you will not be allowed to drive yourself home. No small children are allowed in the surgery center waiting area.

- **6:00AM-** The day prior to your procedure you will begin a CLEAR LIQUID diet. NO SOLID FOODS! Follow the clear liquid diet listed above. You can have as much of the clear liquids that you want throughout the day. We encourage you to stay hydrated with water. ·
- **2:00 PM- Open 1 flavor enhancing packet and put the contents into 1 of the provided bottles.**
- Fill the provided bottle with lukewarm water up to the fill line. After capping the bottle gently shake the bottle until all powder has dissolved. For the best taste, refrigerate the solution for one hour before drinking, Do not freeze.
- **3:00 PM-** Begin drinking the bottle of solution that you have just prepared. **Drink 8 ounces every 15 minutes until the bottle is empty**

If nausea, bloating or abdominal cramping occurs, pause or slow the rate of drinking the solution and drink additional water until the symptoms diminish.

If you choose, this would be a good opportunity to mix your second dose of preparation and refrigerate. You will begin drinking the second dose tomorrow morning.

THE DAY OF YOUR PROCEDURE: DOSE 2

- **3:00AM- Remove from the refrigerator and begin drinking the bottle of solution that you have prepared. Drink 8 ounces every 15 minutes until the bottle is empty**
- **You can drink an additional 16 ounces of water up to FOUR HOURS BEFORE YOUR SCHEDULED TIME FOR YOUR PROCEDURE.**
- **FOUR HOURS BEFORE YOUR PROCEDURE TIME—STOP ALL LIQUIDS. YOU CANNOT HAVE ANYTHING BY MOUTH UNTIL AFTER YOUR PROCEDURE.**

If nausea, bloating or abdominal cramping occurs, pause or slow the rate of drinking the solution and drink additional water until the symptoms diminish.

- If you have medications that you must take in the morning (with the exception of Insulin or blood thinners) you may take them with a **very small sip of water before 5:00AM**
- Wear comfortable clothes and things that are easy to take off and put back on. We recommend wearing shoes that easily slip off and on and don't require you to bend over to put them on.
- Warm socks
- Insurance cards and ID
- Cash or Credit Card if you have a Co-Payment or deductible that has not been met
- A list of all your prescriptions and over the counter medications
- Please avoid wearing contact lenses, jewelry or valuables
- You will be asked to remove any dentures.

PLEASE LEAVE LARGE BAGS AND BACKPACKS AT HOME!

IF YOU TAKE BLOOD THINNERS, STOP 3-7 DAYS BEFORE YOUR APPOINTMENT, FOLLOWING THE DIRECTIONS BELOW:

- **Coumadin, Lovenox, Plavix, Effient, Brillinta, Ticlid, Persatine or any medication that thins your blood please follow up with us as you may need to stop these medications 3-7 days before your procedure.**
- **Lovenox (Enoxaparin) - Hold PM dose the day BEFORE the procedure/ hold AM dose the Day of procedure**
- **Aggrenox (Aspirin-Dipyridamole)- 48 hours prior to procedure**
- **Eliquis (Apaxiban)- 48 hours prior to procedure**
- **Pradaxa (Dabigatran Etexilate) - 48 hours prior to procedure**
- **Xarelto (Rivaroxaban) - 48 Hours prior to procedure**
- **Warfarin (Coumadin)- 3 days prior to procedure**

- Brillianta (Ticagrelor) - 5 days prior to procedure
- Effient (Prasugrel) – 5 days prior to procedure
- Plavix (Clopidogrel)- 5 Days prior to procedure
- Aspirin- you do not need to stop.

IF YOU TAKE DIABETIC MEDICATIONS, FOLLOW THE DIRECTONS BELOW:

- Lantus or Long Acting Insulin: Take only ½ the dose at midnight before your procedure. **Bring your insulin in with you day of procedure**
- Glimperide (Amaryl): Hold day of procedure
- Glucotrol (Glipizide): Hold day of procedure
- Glucophage (Metformin): Hold day of procedure
- Actos (Pioglitazone): Hold day of procedure
- Invokana (Canaglifozin): Hold day of procedure
- Januvia (Sitagliptin): Hold day of procedure

PLEASE MONITOR YOUR BLOOD SUGAR DURING YOUR PREPARATION AND IN THE MORNING OF YOUR PROCEDURE. TAKE DIABETIC MEDICATIONS AS NEEDED. BRING YOUR INSULIN WITH YOU FOR YOUR PROCEDURE.

IF YOU TAKE GLP-1 AGONIST MEDICATIONS FOR TYPE 2 DIABETES OR WEIGHT LOSS, HOLD FOR ONE WEEK PRIOR TO YOUR PROCEDURE. SEE BELOW LIST OF GLP-1 AGONIST MEDICATIONS:

- Ozempic
- Wegovy
- Mounjaro
- Trulicity
- Rybelsus
- Victoza
- Byetta
- Saxenda
- Tirzapetide
- Semaglutide
- Retatrutide