



FIRST SURGERY CENTER

AFFILIATED WITH TRINITY MEDICAL MULTI-SPECIALTY GROUP, INC

COLONOSCOPY GOLYTELY/GAVILYTE PREP

****YOU WILL NEED A PRESCRIPTION FROM THE CLINIC FOR GOLYTELY. This would have been sent to your pharmacy on the day of your clinic visit. If you have questions last minute there may not be time to reach anyone in the surgery center to help address your concerns.**

TIME: You will receive a call from the Surgery Center 3-6 days prior to your appointment with your arrival time.

Please be sure to follow your preparation and all of the instructions detailed below. If you have access to your patient portal, the time you see listed on the schedule is NOT your scheduled time.

ARRANGE FOR A DRIVER: YOU WILL NEED A RESPONSIBLE ADULT TO DRIVE YOU HOME AFTER YOUR PROCEDURE. **You will not be allowed to drive yourself home due to the sedation that you will be given.** Small children are not allowed in the surgery center waiting room and drivers will be asked to wait in their car.

If you find it necessary to change or cancel your appointment, you must do this at least 7 working days prior to your appointment or you will be charged a **\$200** cancellation fee. Please call our office Monday – Friday, 8AM to 4PM at 951-517-0906 if you need to change or cancel your appointment. Follow the prompts for First Surgery Center scheduling.

Your procedure will be performed at:

- () First Surgery Center
31625 De Portola Rd. Suite #102
Temecula, CA 92592
Phone: (951) 517-0906
- () Other _____

THE ITEMS THAT YOU NEED TO PURCHASE AT LEAST 48 HOURS PRIOR TO YOUR PROCEDURE:

- (PRESCRIPTION REQUIRED) –GOLYTELY/GAVILYTE prep solution. If we have not given you a prescription or called it into your pharmacy, please call us at 951-517-0906.
- CLEAR LIQUIDS (SEE SUGGESTIONS BELOW)
- 1 BOTTLE OF SIMETHICONE (a common brand name is Mylicon) (this comes with a dropper for use)

Some examples of clear liquids you are able to have:

Water	Apple Juice or WHITE grape juice	Hard Candy
Broth (Chicken, Vegetable or Bone)	Popsicles (No Red or Purple)	
7up, Ginger ale, Sprite	Jell-O (No Red or Purple)	
Coffee & Tea (No cream)	Sugar or Honey	
	NO DAIRY AND NO ORANGE JUICE	

ONE WEEK BEFORE YOUR PROCEDURE:

- Try to refrain from eating seeds, nuts, corn, popcorn, celery and other high fiber food or supplements
- If you are taking Blood Thinners, GLP-1, or Diabetic Medication, REVIEW AND THEN FOLLOW THE DIRECTIONS ACCORDINGLY ON THE LIST PROVIDED BELOW. If you have questions regarding this, please call our office.

THE DAY BEFORE YOUR PROCEDURE:

YOU WILL NEED A RESPONSIBLE ADULT TO DRIVE YOU HOME AFTER YOUR PROCEDURE. Confirm with your driver, as you will not be allowed to drive yourself home. No small children are allowed in the surgery center waiting area.

- **6:00AM-** The entire day prior to your procedure you will begin a CLEAR LIQUID diet. NO SOLID FOODS! Follow the clear liquid diet listed above. You can have as much of the clear liquids that you want throughout the day. We encourage you to stay hydrated with water. Drink at least 8 glasses of clear liquids throughout the day.
- **8:00AM-** Fill the Golytely jug with cold water and **10 droppers full** of the Simethicone and shake well. Put the jug into the refrigerator. It tends to taste better cold (REFRIGERATION IS OPTIONAL)
- **3:00PM-** Begin drinking the Golytely. Drink 10-11, 8oz cups of Golytely. (It is ok to finish up to midnight). You can drink this solution with a straw if it makes it easier. Sucking on a hard candy may also help you if you dislike the taste of the solution.

****If you experience nausea or vomiting, take a 30 min break and then continue the preparation****

If you have any questions regarding your preparation, please call 951-517-0906

REMINDER: CLEAR LIQUIDS ONLY UNTIL AFTER YOUR PROCEDURE!!!! NO SOLID FOOD!!!! AND STOP ALL FLUID INTAKE NO LESS THAN 4 HOURS BEFORE YOUR CHECK IN TIME.

DAY OF YOUR PROCEDURE:

- **STOP ALL FLUID INTAKE NO LESS THAN 4 HOURS BEFORE YOUR CHECK IN TIME.**
- Take any hypertension, thyroid, anti-seizure, and heart medications as prescribed with a small sip of water. **Avoid taking insulin or blood thinners.** (Unless otherwise advised)
- Leave all large purses, bags and backpacks at home
- Please avoid wearing contact lenses, jewelry or valuables

WHAT TO BRING/DO THE DAY OF YOUR APPOINTMENT:

- Wear comfortable clothes and things that are easy to take off and put back on. We recommend wearing shoes that easily slip off and on and don't require you to bend over to put them on.
- Warm socks
- Insurance cards and ID
- Cash or Credit Card if you have a Co-Payment or deductible that has not been met
- A list of all your prescriptions and over the counter medications

DO NOT BRING LARGE BAGS OR BACKPACKS TO YOUR APPOINTMENT

IF YOU TAKE BLOOD THINNERS, STOP 3-7 DAYS BEFORE YOUR APPOINTMENT, FOLLOWING THE DIRECTIONS BELOW:

- Coumadin, Lovenox, Plavix, Effient, Brillinta, Ticlid, Persatine or any medication that thins your blood please follow up with us as you may need to stop these medications 3-7 days before your procedure.
- Lovenox (**Enoxaparin**) - Hold PM dose the day BEFORE the procedure/ hold AM dose the Day of procedure
- Aggrenox (**Aspirin-Dipyridamole**)- 48 hours prior to procedure
- Eliquis (Apaxiban)- 48 hours prior to procedure

- Pradaxa (**Dabigatran Etexilate**) - 48 hours prior to procedure
- Xarelto (**Rivaroxaban**) - 48 Hours prior to procedure
- Warfarin (Coumadin)- 3 days prior to procedure
- Brillianta (**Ticagrelor**) - 5 days prior to procedure
- Effient (Prasugrel) – 7 days prior to procedure
- Plavix (Clopidogrel)- 7 Days prior to procedure
- Aspirin- you do not need to stop.

IF YOU TAKE DIABETIC MEDICATIONS, FOLLOW THE DIRECTONS BELOW:

- Lantus or Long Acting Insulin: Take only ½ the dose at midnight before your procedure. Bring your insulin in with you day of procedure
- Glimperide (Amaryl): Hold day of procedure
- Glucotrol (Glipizide): Hold day of procedure
- Glucophage (Metformin): Hold day of procedure
- Actos (Pioglitazone): Hold day of procedure
- Invokana (Canaglifozin): Hold day of procedure
- Januvia (Sitagliptin): Hold day of procedure

IF YOU TAKE GLP-1 AGONIST MEDICATIONS FOR TYPE 2 DIABETES OR WEIGHT LOSS, HOLD FOR ONE WEEK PRIOR TO YOUR PROCEDURE. SEE BELOW LIST OF GLP-1 AGONIST MEDICATIONS:

- **Ozempic**
- **Wegovy**
- **Mounjaro**
- **Trulicity**
- **Rybelsus**
- **Victoza**
- **Byetta**
- **Saxenda**
- **Tirzapetide**
- **Semaglutide**
- **Retatrutide**